

## A Biochemical, Histologic, and Immunohistologic Analysis of Membranes Obtained From Failed Cemented and Cementless Total Knee Arthroplasty

JUNJI CHIBA, M.D.,\* LESLIE J. SCHWENDEMAN, M.D.,\*\* ROBERT E. BOOTH, JR., M.D.,†  
LAWRENCE S. CROSSETT, M.D.,\*\* AND HARRY E. RUBASH, M.D.\*\*

Biochemical, histologic, and immunohistochemical analyses were performed on 34 interface membranes obtained from 33 patients during revision total knee arthroplasty. The membranes had surrounded components of cementless (n = 11) and cemented (n = 23) knee prostheses that were aseptically loose. None of these implant failures was caused by catastrophic polyethylene erosion leading to metal-to-metal contact. The histologic findings were similar in the membranes from cemented and cementless knee components: small polyethylene debris within macrophages and large birefringent polyethylene debris within foreign-body giant cells. Metallic debris was seen in membranes from both groups, but cemented membranes had more polymethylmethacrylate particles and more hyalinization. Intracytoplasmic asteroid bodies were observed in several foreign-body giant cells in both types of membranes. No significant differences were found between the two groups in levels of collagenase, prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), interleukin-1 (IL-1), interleukin-6 (IL-6), or tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), nor in the population of inflammatory cells stained with IL-1, IL-6, and TNF- $\alpha$  antibodies. Membranes that had surrounded components with radiographic evidence of

diffuse or localized periprosthetic bone loss released significantly more collagenase, IL-1, IL-6, and TNF than did membranes from components without bone loss. These two groups, however, did not have significantly different PGE<sub>2</sub> levels. These findings suggest that polyethylene and metal debris may play a role in macrophage activation and the release of mediators of bone resorption in the membranes surrounding failed cemented and cementless total knee implants.

Aseptic loosening and osteolysis have increased as the premier obstacle limiting the longevity of total joint replacements. Initially thought to be inherent to the use of polymethylmethacrylate (PMMA) cement, osteolysis is now recognized in both cemented and uncemented arthroplasties. Its occurrence is well documented in total hip arthroplasty, and is believed to be the result of a foreign body reaction to particulate PMMA, metal, and polyethylene wear debris.<sup>7,11,13,15,17-20,23,28</sup> There is also evidence that several biochemical factors associated with bone resorption such as collagenase, prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), interleukin-1 (IL-1), interleukin-6 (IL-6), and tumor necrosis factor (TNF), may play an important role in the development of aseptic loosening and osteolysis.<sup>5,7,15,17</sup>

The problem of osteolysis has not been as well described in total knee arthroplasty (TKA). Recently, however, investigators have noted osteolysis around loosened cementless total knee prostheses.<sup>14,21</sup> These re-

ports have described implant interface histology to the osteolysis around cemented and uncemented total knee prostheses. Biochemical analysis of membranes obtained from total knee arthroplasties has been documented, but no systematic study was conducted. This study was a histologic, and immunohistologic analysis of the characteristics of interface membranes from failed cemented and uncemented total knee arthroplasties (referred to as "cemented" and "cementless" total knee arthroplasties).

### MATERIALS AND METHODS

Thirty-four interface membranes were obtained from 33 patients who had undergone revision failed total knee arthroplasty at the University of Pittsburgh Medical Center. The patients were divided into two groups: cemented (n = 23) and cementless (n = 11). Twenty-three membranes were obtained from cemented total knee arthroplasties, ten Insall and five Johnson. The mean age was 65 years. Eleven cementless membranes were obtained from Miller-Galante total knee arthroplasties. The seven patients had a mean age of 68 years. To serve as a control, ten membranes were obtained from six cemented and four cementless total knee arthroplasties. The pseudocapsule of the revision knee was removed from all three groups and the patellar and tibial components were removed with carbon-implant.

The average time from total knee arthroplasty and revision was 7.1 years (range, 49-109 months). The average time from total knee arthroplasty and revision was 80.6 months for the cementless group. The first arthroplasty was cemented in eight patients and eight cementless. The indications for revision were knee pain and instability in ten cemented and eight cementless patients. The amount of bone loss was defined, on preoperative radiographs, as the distance between the radiolucent line at the cement or bone-implant interface and the radiolucent line at the bone-implant interface.

\* The Institute of Rheumatology, Tokyo Women's Medical College, Nishi-shinjuku, Shinjuku-ku, Tokyo, Japan.

\*\* Ferguson Laboratory, Department of Orthopaedic Surgery, University of Pittsburgh, Pittsburgh, Pennsylvania.

\*\* The Rothman Institute, Pennsylvania Hospital, Philadelphia, Pennsylvania.

Reprint requests to Junji Chiba, M.D., c/o Harry E. Rubash, M.D., 3471 Fifth Ave., Liliane Kaufmann Building, Suite 1010, Pittsburgh, PA 15213.

Received and accepted: May 3, 1993.

# Immunohistologic From Failed Total Knee Arthroplasty

† E. BOOTH, JR., M.D.,†  
BASH, M.D.\*\*

Periprosthetic bone loss requires more collagenase, IL-1, IL-6, and membranes from components of these two groups, however, did not differ PGE<sub>2</sub> levels. These two polyethylene and metal debris in macrophage activation and bone resorption in the failed cemented and cementless implants.

Loosening and osteolysis have been the primary obstacle limiting the long-term success of total knee replacements. Initially attributed to the use of polyethylene (PMMA) cement, osteolysis occurred in both cemented and uncemented prostheses. Its occurrence is not unique to total hip arthroplasty, but may be the result of a foreign body reaction to particulate PMMA, metal, or wear debris.<sup>7,11,13,15,17-20,23,28</sup> Evidence that several biochemical mediators, including prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), IL-1, interleukin-6 (IL-6), and tumor necrosis factor (TNF), may play an important role in the development of aseptic loosening.<sup>5,7,15,17</sup>

Although osteolysis has not been as well studied in total knee arthroplasty as in hip, however, investigators have reported osteolysis around loosened cemented prostheses.<sup>14,21</sup> These re-

ports have described the presence of bone-implant interface membranes with similar histology to those reported in cases of osteolysis around cementless total hip prostheses. Biochemical analyses of these membranes from total knee replacement have not been documented, however. The current study was conducted to determine the biochemical, histologic, and immunohistochemical characteristics of interface membranes surrounding failed cementless and cemented TKAs (referred to as "cementless membranes" and "cemented membranes", respectively).

## MATERIALS AND METHODS

Thirty-four interface membranes were harvested from 33 patients undergoing surgery to revise failed total knee prostheses at the University of Pittsburgh Medical Center and the Pennsylvania Hospital between July 1991 and April 1992. Twenty-three membranes (22 patients) were taken from cemented implants: five porous-coated anatomic, ten Insall-Burstein, three Miller-Galante, and five Johnson & Johnson prostheses. This group comprised 12 men and 11 women whose mean age was 65.7 years (range, 47-81 years). The 11 cementless membranes were taken from nine Miller-Galante and two LCS Rotating Platform implants. The seven women and four men in this group had a mean age of 63.5 years (range, 38-83 years). To serve as a control group for the biochemical analysis, ten membranes (four cementless and six cemented) were harvested from the fibrous pseudocapsule of ten additional patients undergoing revision knee arthroplasty. All components in all three groups articulated with a polyethylene tibial and patellar surface. There were no patients with carbon-impregnated polyethylene.

The average time between the initial arthroplasty and revision procedure was 60.1 months (range, 49-109 months) in the cemented group and 80.6 months (range, 26-96 months) in the cementless group. The primary indications for the first arthroplasty were osteoarthritis (13 cemented and eight cementless) and rheumatoid arthritis (ten cemented and three cementless), whereas the indications to perform all revision procedures were knee pain and radiographic findings of component loosening or migration. The latter was defined, on preoperative radiographs, in accordance with the Knee Society system as the presence of a radiolucent line at least 1 mm wide at the bone-cement or bone-implant interface (Fig. 1).<sup>6,26</sup> The widths of radiolucent lines were measured on the

basis of seven zones on an anteroposterior view of the tibia; three on a lateral view of the tibia and seven on a lateral view of the femur. The widths of radiolucent lines on a tangential radiograph of the patella were not measured. Among the 23 cemented implants, eight femoral components and 13 tibial components met these criteria; in the 11 cementless implants, the numbers were three femoral and six tibial components (Table 1). Intraoperative examination confirmed that all of these components were loose and surrounded by a fibrous membrane. Loosening was confirmed intraoperatively by using manual pressure to elicit visible motion at the bone-implant interface.

All patients in this study had discontinued taking antiinflammatory medications ten days before revision surgery. In addition, preoperative and intraoperative cultures of aspirates from the hip joint were sterile for all patients.

All membranes were cut into portions for separate biochemical, histologic, and immunohistochemical analysis.

## HISTOLOGY

Periprosthetic membrane samples were prepared for hematoxylin and eosin (H&E) staining. Paraffin-embedded specimens were analyzed qualitatively under regular light and polarized microscopy for macrophages, fibroblasts, particulate PMMA, metal debris, and polyethylene debris. Using a modification of Mirra's grading system,<sup>15,20</sup> each sample was graded according to the number of cells and particles present (0 = absent, 1+ = mild, 2+ = moderate, 3+ = severe).

## BIOCHEMICAL ASSAYS

The membrane specimens were cut into small fragments, and organ cultures were performed using a modified Goldring method.<sup>6</sup> After three days of incubation, the culture media were recovered and assayed for collagenase, PGE<sub>2</sub>, IL-1, IL-6, and TNF- $\alpha$ .

Collagenase activity was measured using post-synthetically labeled [<sup>3</sup>H] acetic anhydride (New England Nuclear, Boston, Massachusetts) as a substrate. The detailed technique was reported previously.<sup>15,27</sup> One unit of collagenase activity degrades 1  $\mu$ g per minute of substrate. Prostaglandin E<sub>2</sub> was measured by radioimmunoassay with a commercially available kit (New England Nuclear), and IL-1 was measured by bioassay using the proliferation of the D10.G4.1 T helper clone, as described elsewhere.<sup>15,25</sup> Finally, IL-6 and TNF- $\alpha$  were measured with a commercially available enzyme-linked immunosorbent assay (ELISA) kit.




FIG. 1. Anteroposterior and lateral radiograph of a loosened cementless TKA.

The data thus obtained were statistically compared with a Student's *t*-test.

#### IMMUNOHISTOCHEMISTRY

To clarify which cells were producing IL-1, IL-6, and TNF- $\alpha$  in the interface membranes, immunohistochemical analysis was performed using anti-IL-1, anti-IL-6, and anti-TNF- $\alpha$  antibodies. The specimens in this study were frozen in OCT compound (Lab-Tek Products, Naperville, Illinois) and stored at  $-70^{\circ}$ . Sections of the frozen tissue were cut 7  $\mu$ m thick on a cryostat, fixed in cold acetone ( $4^{\circ}$ ) for ten minutes, and then dried. After these sections were washed in phosphate-buffered saline (PBS) for five minutes, they were incubated for ten minutes in an avidin- and biotin-blocking reagent (Vector Laboratory, Burlingame, California). They were washed again in PBS for five minutes and incubated with normal horse

serum for 30 minutes at room temperature with monoclonal antibodies to macrophages, B lymphocytes (anti-CD3), and T lymphocytes (anti-CD22) (DAKOPATTS, Glostrup, Denmark) and polyclonal antibodies to IL-1, IL-6, and TNF- $\alpha$  (Calbiochem, La Jolla, California). Endogenous peroxidase was neutralized with 0.05% hydrogen peroxide for ten minutes. Next the biotinylated secondary antibody (Vector Laboratory) was applied to the sections for 45 minutes at room temperature. Then they were washed with PBS, after which 3,3'-diaminobenzidine substrate (Sigma, St. Louis, Missouri) with 0.05% hydrogen peroxide was added for color development. Finally, the sections were rinsed for five minutes, exposed to hematoxylin, and mounted with Immunomount (Shandon, Sewickley, Pennsylvania). Control specimens had the same preparation except that the first antibodies were omitted.

TA

Zone

1  
2  
3  
4  
5  
6  
7

RADIOG

Eight of  
nents and  
component  
1 mm at t  
larly, three  
ponents an  
component  
ria (Table  
confirmed  
and surrou

In both  
membranes

FIG. 2. L  
specimen tak  
around the ti  
strates partic  
is limited adja  
faces.

TABLE 1. Percentage of Knees With Radiolucent Lines According to Zone

Zone	Radiographs					
	Lateral of Femur		Anteroposterior of Tibia		Lateral of Tibia	
	Cemented	Cementless	Cemented	Cementless	Cemented	Cementless
1	5	3	6	3	2	1
2	3	0	4	3	1	1
3	0	0	2	0	0	0
4	0	0	1	0		
5	0	0	0	0		
6	0	0	0	0		
7	0	0	0	0		

## RESULTS

### RADIOGRAPHIC AND INTRAOPERATIVE FINDINGS

Eight of the 23 cemented femoral components and 13 of the 23 cemented femoral components had a radiolucent line of at least 1 mm at the bone-cement interface. Similarly, three of the 11 cementless femoral components and six of the 11 cementless tibial components were loose by radiographic criteria (Table 1). Intraoperative examination confirmed that these components were loose and surrounded by a fibrous membrane.

### HISTOLOGIC FINDINGS

In both the cemented and cementless membranes, microscopic examination showed

macrophage infiltrates, multinucleated foreign-body giant cells, and large accumulation of debris (metal, polyethylene, or cement) within dense, organized connective tissue stroma. Metal debris was seen in membranes from both experimental groups, and most of it was intracellular within macrophages. In addition, the particulate metal frequently was confined to the membrane surface adjacent to the implant (Fig. 2). Particulate polyethylene was universally present in membranes from both groups. Many small polyethylene particles were contained within the cytoplasm of macrophages; larger particles were engulfed by foreign-body giant cells. Membranes from cemented prostheses generally contained PMMA particles and increased tissue hyalinization. Furthermore, most of the polyethylene particles were lo-

FIG. 2. Light microscopy of the specimen taken from Zones 1 and 2 around the tibial component demonstrates particulate metal debris, which is limited adjacent to the implant surfaces.



entless TKA.

at room temperature with to macrophages, B lymphocytes (anti-CD4, Glostrup, Denmark) and CD4, IL-1, IL-6, and TNF- $\alpha$  (Calbiochem, La Jolla, California). Endogenous peroxidase was blocked with 0.05% hydrogen peroxide for 30 minutes. Next the biotinylated secondary antibody (Vector Laboratories) was applied for 45 minutes at room temperature, washed with PBS, after avidin substrate (Sigma, St. Louis, MO). 0.05% hydrogen peroxide was added for 30 minutes. Finally, the section was exposed to heated with Immunomount (Shandon Scientific Company, Pittsburgh, Pennsylvania). Control specimen was prepared except that the tissue was not fixed.

cated within the specimen where it faced the bone. Interestingly, the large particles of birefringent polyethylene appeared to occupy the periphery of the membrane. Several of the foreign-body giant cells in both types of membranes also contained intracytoplasmic asteroid bodies (Fig. 3). The control tissues were composed of a dense collagenous stroma containing no wear particles and few inflammatory cells.

#### BIOCHEMICAL FINDINGS

The collagenase activity in culture media recovered from the cementless membranes ranged from 1.8 to 14.4 units per gram of tissue with a mean value ( $\pm$  standard error of the mean [SEM]) of  $5.93 \pm 5.17$ . For the cemented membranes, the range of collagenase activity was 1.2–13.7 units per gram of tissue, with a mean of  $5.79 \pm 4.47$ . The values for both experimental groups were not significantly different. Both sets of values were significantly higher ( $p < 0.01$ ) than those for control tissues, however, which ranged from 0 to 7.3 units per gram of tissue, with a mean of  $2.3 \pm 1.2$  (Fig. 4A).

Likewise, the amounts of  $PGE_2$  in cemented membranes (range, 0–13.9 ng/mg of




FIG. 3. Asteroid bodies (short arrows) and large pieces of birefringent polyethylene debris (long arrows) are visible within foreign-body giant cells. At least two asteroid bodies can be seen in this plane of focus. (Stain, hematoxylin and eosin; original magnification,  $\times 400$ .)

tissue; mean,  $5.52 \pm 3.75$ ) and cementless membranes (range, 1.4–14.3 ng/mg of tissue; mean,  $4.94 \pm 3.83$ ) were not significantly different. Again, both sets of values were significantly higher ( $p < 0.05$ ) than control tissue values (range, 0–3.01 ng/mg tissue; mean,  $1.93 \pm 1.01$ ) (Fig. 4A).



The activity of IL-1 was measurable in six of the 11 cementless membranes (range, 0–3.9 units per 100 mg of tissue; mean,  $0.52 \pm 0.57$ ) and in 11 of the 23 cemented membranes (range, 0–0.66 units per 100 mg tissue; mean,  $0.39 \pm 0.82$ ). There was no significant difference between these two groups, but both produced significantly higher IL-1 activity than did control tissues ( $p < 0.05$ ), in which activities were uniformly undetectable (Fig. 4B).

Interleukin-6 activity was detected in all 11 cementless membranes (range, 0–303.5 ng/g tissue; mean,  $123.1 \pm 108.4$ ) and all 23 cemented membranes (range, 0–311.7 ng/g of tissue; mean,  $102.6 \pm 103.1$ ). Tumor necrosis factor- $\alpha$  activity was detected in eight cementless membranes (range, 0–5.9 ng/g tissue; mean,  $1.63 \pm 1.48$ ) and 19 cemented membranes (range, 0–9.4 ng/g tissue; mean,  $2.03 \pm 3.29$ ). There was also no significant difference between these two groups in either factor, and again, both groups had higher levels of both factors as compared with control specimens (Fig. 4B and 4C).

Table 2 compares the activity levels of the biochemical factors as measured in culture media recovered from the membranes. Levels of collagenase,  $PGE_2$ , IL-1, IL-6, and TNF- $\alpha$  did not differ significantly between experimental groups, but both of these groups had significantly higher levels of all of these than did the control group.

Both linear and lytic bone loss were found in association with failed femoral and tibial implants, regardless of whether the prosthesis was cemented. Postoperative radiographs showed that the medial aspect of the tibial metaphysis (Zones 1 and 2) was the most common site for osteolytic bone resorption. Intraoperatively, all of these implants were

Number 2  
February



FIGS. 4  
branes, c  
cemente  
in each g  
circles.

found to  
osteolys  
membra  
tained c  
had phag  
bris (Fig

TA

Collagenase  
(units/g)  
 $PGE_2^*$  (ng/  
IL-1\* (u/10  
IL-6\* (ng/g  
TNF- $\alpha^*$  (ng

n represent  
\* Significan


3.75) and cementless  
1-14.3 ng/mg of tissue;  
re not significantly dif-  
s of values were signifi-  
5) than control tissue  
ng/mg tissue; mean,

was measurable in six  
s membranes (range,  
g of tissue; mean, 0.52  
he 23 cemented mem-  
units per 100 mg tissue;  
here was no significant  
hese two groups, but  
antly higher IL-1 activ-  
tissues ( $p < 0.05$ ), in  
uniformly undetectable

ty was detected in all 11 es (range, 0–303.5 ng/g  $\pm$  108.4) and all 23 ce- range, 0–311.7 ng/g of 103.1). Tumor necrosis was detected in eight ce- (range, 0–5.9 ng/g tis- .48) and 19 cemented –9.4 ng/g tissue; mean, was also no significant ese two groups in either h groups had higher lev- compared with control nd 4C).

the activity levels of the as measured in culture n the membranes. Lev- PGE<sub>2</sub>, IL-1, IL-6, and significantly between ex- out both of these groups her levels of all of these group.

tic bone loss were found. Failed femoral and tibial of whether the prosthesis postoperative radiographs distal aspect of the tibial 1 and 2) was the most osteolytic bone resorption. of these implants were



FIGS. 4A-4C. The results of biochemical comparison among cemented membranes, cementless membranes, and control tissues, represented as amounts produced per gram of membrane. Closed circles, cemented membranes; open circles, cementless membranes. The mean value and standard deviation (SD) in each group are indicated, respectively, by a vertical bar with a long and short horizontal bar beside the circles.

found to be loose. Four cases of focal tibial osteolysis were noted on radiographs. The membranes around these four implants contained concentrations of macrophages that had phagocytized polyethylene and metal debris (Fig. 5). Biochemical results from speci-

mens with radiographic evidence of osteolysis were also regrouped and compared with those specimens without evidence of osteolysis. The specimens from patients with evidence of osteolysis had released significantly higher ( $p < 0.05$ ) amounts of collagenase. IL-

**TABLE 2. Levels of Biochemical Factors in Experimental and Control Groups**

	Cemented			Cementless			Control		
	Mean $\pm$ SEM (range)		n	Mean $\pm$ SEM (range)		n	Mean $\pm$ SEM (range)		n
Collagenase* (units/g)	5.79 $\pm$	4.47 (1.2-13.7)	23	5.93 $\pm$	5.17 (1.8-14.4)	11	2.3 $\pm$ 1.2 (0-7.3)		10
PGE <sub>2</sub> * (ng/mg)	5.52 $\pm$	3.75 (0-13.9)	23	4.94 $\pm$	3.83 (1.4-14.3)	11	1.93 $\pm$ 1.01 (0-3.01)		10
IL-1* (u/100mg)	0.39 $\pm$	0.82 (0-0.66)	11	0.52 $\pm$	0.57 (0-3.9)	6	undetectable		10
IL-6* (ng/g)	102.6 $\pm$	103.1 (0-311.7)	23	123.1 $\pm$	108.4 (0-303.5)	11	38.3 $\pm$ 35.7		10
TNF- $\alpha$ * (ng/g)	2.03 $\pm$	3.29 (0-9.4)	19	1.63 $\pm$	1.48 (0-5.9)	8	0.07 $\pm$ 0.45		10

*n* represents number of specimens in which activity was detected.

\* Significantly higher than control values ( $p < 0.05$ ).



FIG. 5. Polarized microscopy of a specimen taken from Zones 1 and 2 demonstrates birefringent debris within multinucleated foreign-body giant cells and mononuclear histiocytes. (Stain, hematoxylin and eosin; original magnification,  $\times 100$ .)

1, IL-6, and TNF- $\alpha$  into the culture media than did the membranes from prostheses without focal osteolysis (Table 3). There were no statistically significant differences between the levels of PGE<sub>2</sub> in groups with and without osteolysis, however.

#### IMMUNOHISTOCHEMICAL FINDINGS

In the 34 membranes analyzed, numerous macrophages were stained with antimacrophage antibody and were located beneath the synovial cell lining or fibrin layer, which was adjacent to the implant surface (Fig. 6). In sections of all membranes from both experimental groups that were stained sequentially

with anti-CD3 and anti-CD22 monoclonal antibodies, very few T lymphocytes and B lymphocytes, respectively, were evident. Many macrophages (Fig. 7), relatively large numbers of fibroblasts, and some endothelial cells were stained with anti-IL-1, IL-6, and TNF- $\alpha$  antibodies. There were no significant differences between the numbers of anti-IL-6-positive cells in cemented and cementless membranes. In contrast, the populations of macrophages, fibroblasts, and endothelial cells that stained with anti-IL-1, IL-6, and TNF- $\alpha$  antibodies were larger in specimens from patients with bone lysis than in those without this problem.

TABLE 3. Comparison of Chemical Mediators in Condition Media From the Membrane With and Without Osteolysis in Failed Cemented and Cementless Knee Prostheses

	Focal Osteolysis			
	Positive		Negative	
	Cemented (n = 21)	Cementless (n = 9)	Cemented (n = 25)	Cementless (n = 13)
Collagenase (unit/g)	6.82 $\pm$ 5.11*	7.00 $\pm$ 4.46*	4.92 $\pm$ 5.02	5.18 $\pm$ 4.62
PGE2 (ng/g)	5.86 $\pm$ 3.28	5.21 $\pm$ 3.22	5.22 $\pm$ 3.05	4.76 $\pm$ 3.73
IL-1 (U/g)	6.01 $\pm$ 3.85*	6.70 $\pm$ 2.57*	2.11 $\pm$ 6.84	3.70 $\pm$ 5.21
IL-6 (ng/g)	125.5 $\pm$ 86.3*	152.5 $\pm$ 92.6*	83.3 $\pm$ 96.1	102.7 $\pm$ 98.5
TNF- $\alpha$ (ng/g)	3.03 $\pm$ 4.54*	2.91 $\pm$ 4.21*	1.19 $\pm$ 2.02	0.75 $\pm$ 1.26

\*  $p < 0.05$ ; values are the mean  $\pm$  SEM.

FIG. 6. The specimen from a patient with focal osteolysis is adjacent to the implant surface. (Stain, hematoxylin and eosin; original magnification,  $\times 100$ .)

FIG. 7. Staining of a specimen from a patient with focal osteolysis using anti-macrophage antibody. Numerous macrophages are present in the center of the specimen. (Stain, hematoxylin and eosin; original magnification,  $\times 200$ .)

olarized microscopy of a  
ken from Zones 1 and 2  
as birefringent debris  
tinucleated foreign-body  
and mononuclear histo-  
t, hematoxylin and eosin;  
gnification,  $\times 100$ .)

anti-CD22 monoclonal  
T lymphocytes and B  
tively, were evident.  
Fig. 7), relatively large  
s, and some endothelial  
th anti-IL-1, IL-6, and  
ere were no significant  
re numbers of anti-IL-  
mented and cementless  
ast, the populations of  
lasts, and endothelial  
h anti-IL-1, IL-6, and  
re larger in specimens  
one lysis than in those

a From the Membrane  
s Knee Prostheses

Negative	
ed 5)	Cementless (n = 13)
5.02	5.18 $\pm$ 4.62
3.05	4.76 $\pm$ 3.73
6.84	3.70 $\pm$ 5.21
6.1	102.7 $\pm$ 98.5
2.02	0.75 $\pm$ 1.26




FIG. 6. Antimacrophage antibody staining of the specimen shown in Figure 2 demonstrates numerous macrophages beneath a fibrin layer, which is adjacent to the implant surface. (The implant surface is the upper part of the specimen.) (Original magnification,  $\times 200$ .)

## DISCUSSION

Late loosening of total joint prostheses has been considered primarily a biomechanical process. It has become increasingly apparent, however, that periprosthetic tissues and their production of a number of biochemical substances play an important role in aseptic loosening. These interface membranes contribute to the bone resorption and osteolysis associated with aseptic loosening. This process has been previously described in total hip arthroplasty and can occur with cemented or uncemented prostheses and with stable and unstable implants.<sup>7,11,13,15,17,18-20,23,28</sup>

Recently, Nolan and Bucknill<sup>21</sup> and Peters *et al.*<sup>22</sup> investigated osteolysis associated with total knee prostheses. Their histologic findings were similar to those previously found with osteolysis surrounding cementless hip prostheses. They noted submicronic particulate polyethylene or metal debris within macrophages and large pieces of polyethylene debris or particulate PMMA within foreign-body giant cells. The authors also observed large intracytoplasmic asteroid bodies within foreign-body giant cells, however. These giant cells form from a syncytium of macrophages in response to particulate debris too large to be phagocytized by a single cell. As-

FIG. 7. Immunohistochemical staining of the membrane obtained from a patient with focal osteolysis using anti-TNF- $\alpha$ -positive antibody. Numerous anti-TNF- $\alpha$ -positive macrophages are visible. A large fragment of polyethylene is shown in the center of the field. (Original magnification,  $\times 200$ .)





teroid bodies are composed of microtubules and intermediate fibers, two constituents of the normal cellular cytoskeleton.<sup>1,2</sup> Because asteroid bodies were seen in giant cells containing large particles of polyethylene debris, it is speculated that these particles create a microenvironment in which the giant cell cytoplasm undergoes various transformations from solution to a gelatinous form. These sol-gel transformations result in a condensation of the giant cell's cytoskeleton, which results in the formation of the stellate-shaped asteroid bodies. The presence of asteroid bodies may represent a burned out or "aged" population of giant cells.

Interestingly, large fragments of polyethylene, foreign-body giant cells, and fat were more common in the membranes obtained from failed total knee implants than in those from failed total hip implants, regardless of whether they were cemented or cementless. This may be due to the high contact stresses in knee implants and also may reflect the different motion of these implants. Knee implants have a very complicated motion, consisting of rolling, gliding, and sliding. In contrast, hip implants predominantly glide, producing small powderlike polyethylene and metal debris, which can stimulate macrophages, thus inducing a more rapid and extensive osteolysis than occurs in the knee. A different histologic and biochemical response in the interface membranes from failed cemented and cementless knee prostheses may explain the lower level of progressive osteolysis seen with these implants.

It was impossible to estimate how many particles each macrophage had phagocytized because of their irregularity, and because some macrophages were spread on the surface of the large particles. In addition, recent reports have indicated that many of these particles were too small to be resolved with light microscopy; therefore, microscopic analysis may not provide an accurate estimate of the amount of wear debris in the membranes.

Kim *et al.*<sup>15</sup> found substantial levels of collagenase, PGE<sub>2</sub>, and IL-1 in the membranes

from failed cementless femoral components in total hip arthroplasty—levels not significantly different from those seen in the membranes from failed cemented femoral components.<sup>15</sup> Likewise, no significant differences were found between cemented and cementless knee membranes in the levels of collagenase, PGE<sub>2</sub>, IL-1, IL-6, and TNF- $\alpha$ . Nonetheless, in the current study, collagenase, IL-1, IL-6, and TNF- $\alpha$  levels were significantly higher in specimens from patients with focal osteolysis. The medial aspect of the tibial metaphysis was the most common site for osteolytic bone resorption, consistent with findings by Peters *et al.*<sup>22</sup> Although biochemical agents such as collagenase, PGE<sub>2</sub>, IL-1, IL-6, and TNF- $\alpha$  are associated with bone resorption,<sup>3,5,7,8,12,16</sup> cytokines such as bone morphogenic protein and transforming growth factor- $\beta$  should have been measured. Their activity may appear to be inhibited, because these cytokines are believed to stimulate osteoblasts.

It is well known that several biochemical mediators, including various cytokines, contribute to bone resorption.<sup>9,10</sup> It is unclear why membranes from patients with linear or lytic bone loss and those without bone loss did not contain significantly different levels of PGE<sub>2</sub>, given that bone loss was associated with higher levels of all other biochemical factors measured. Interleukin-1 and TNF- $\alpha$  not only induce the production of IL-6 by macrophages, fibroblasts, and endothelial cells, but they also induce the proliferation of fibroblasts and endothelial cells.<sup>24</sup> Furthermore, IL-1 and TNF- $\alpha$  stimulate osteoclastic bone resorption. It was recently reported that IL-6 appears to induce osteoclastic bone resorption directly and to inhibit the proliferation of fibroblasts.<sup>12,16,24</sup> The biochemical and immunohistochemical results reported here suggest that macrophages, fibroblasts, and endothelial cells in the membranes react with metal or polyethylene debris and produce IL-6, which directly stimulates osteoclasts in patients with linear or lytic bone loss. Because IL-1 and TNF- $\alpha$  both stimulate os-

teod  
6,<sup>9,16</sup>  
direc  
duct  
fibro  
likel  
cont  
beca  
serve  
tions  
tivity  
chron  
throu  
spons  
Th  
abilit  
mem  
the c  
sorpti  
in the  
mente  
After  
variou  
surrou  
tion b  
polyet  
surface  
thetic  
lease o  
ethyle  
terface  
foreign  
Activa  
such as  
the me  
a num  
with bo  
particu  
macrop  
IL-1, IL  
import  
mecha  
play a  
theses.

The au  
sistance is

ntless femoral components  
oplasty—levels not signifi-  
om those seen in the mem-  
l cemented femoral compo-  
, no significant differences  
en cemented and cement-  
ines in the levels of collage-  
IL-6, and TNF- $\alpha$ . Nonethe-  
nt study, collagenase, IL-1,  
 $\alpha$  levels were significantly  
ns from patients with focal  
edial aspect of the tibial me-  
nost common site for osteo-  
tion, consistent with find-  
al.<sup>22</sup> Although biochemical  
llagenase, PGE<sub>2</sub>, IL-1, IL-6,  
ssociated with bone resorp-  
okines such as bone mor-  
and transforming growth  
ave been measured. Their  
ear to be inhibited, because  
re believed to stimulate os-

n that several biochemical  
ling various cytokines, con-  
resorption.<sup>9,10</sup> It is unclear  
from patients with linear or  
nd those without bone loss  
significantly different levels  
at bone loss was associated  
ls of all other biochemical  
l. Interleukin-1 and TNF- $\alpha$   
the production of IL-6 by  
broblasts, and endothelial  
o induce the proliferation of  
ndothelial cells.<sup>24</sup> Further-  
TNF- $\alpha$  stimulate osteoclastic  
It was recently reported that  
nduce osteoclastic bone re-  
and to inhibit the prolifera-  
sts.<sup>12,16,24</sup> The biochemical  
tochemical results reported  
t macrophages, fibroblasts,  
ells in the membranes react  
olyethylene debris and pro-  
h directly stimulates osteo-  
with linear or lytic bone loss.  
d TNF- $\alpha$  both stimulate os-

teoclastic bone resorption<sup>8,24</sup> and induce IL-  
6,<sup>9,10,12</sup> IL-1 and TNF- $\alpha$  also may contribute  
directly as well as indirectly, through the in-  
duction of IL-6 production by macrophages,  
fibroblasts, and endothelial cells. It is un-  
likely that B lymphocytes and T lymphocytes  
contribute significantly to IL-6 production,  
because these two cell types were rarely ob-  
served in the current study. These observa-  
tions suggest that significant levels of IL-6 ac-  
tivity were produced through a nonspecific  
chronic inflammatory reaction rather than  
through a lymphocyte-mediated immune re-  
sponse.

The results of this study demonstrate the  
ability of cementless and cemented knee  
membranes to produce similar amounts of  
the chemical agents associated with bone re-  
sorption. These factors may play a major role  
in the process of aseptic loosening of both ce-  
mented or cementless total knee prostheses.  
After the implantation of a prosthesis,  
various biochemical factors influence the  
surrounding cells. Abrasion of the articula-  
tion between the femoral component and the  
polyethylene articulating tibial and patellar  
surface as well as the micromotion of the pro-  
sthetic stem may result in the continual re-  
lease of metal, particulate PMMA, and poly-  
ethylene debris. Subsequently, activated in-  
terface membranes can be produced by the  
foreign-body reaction initiated by this debris.  
Activation of various inflammatory cells  
such as macrophages and fibroblasts within  
the membrane can induce the production of  
a number of cytokines that are associated  
with bone resorption. The phagocytization of  
particulate polyethylene and metal debris by  
macrophages can induce the production of  
IL-1, IL-6, and TNF- $\alpha$ , which are considered  
important factors in bone resorption. This  
mechanism of aseptic loosening appears to  
play a role in failed knee as well as hip pros-  
theses.

#### ACKNOWLEDGMENTS

The authors thank Keith Short, B.S., for technical as-  
sistance in the preparation of histologic specimens; Lori

McIntyre Larkin, B.S., for IL-1 assays; Y. Iwaki, M.D.  
(Department of Immunology, University of Pittsburgh),  
for his cooperation with the immunohistochemical anal-  
ysis; and Chris Evans, Ph.D., for his helpful discussions.  
They also thank Jim Dowd, M.D., and Helene Marion  
for their editorial assistance, and Robin Evans, R.N., for  
the preparation of the manuscript.

#### REFERENCES

1. Cain, H., and Kraus, B.: The ultrastructure and morphogenesis of asteroid bodies in sarcoidosis and other granulomatous disorders. *In* Williams, W. W., and Davies, B. H. (eds.): Sarcoidosis, Cardiff, Alpha Omega Press, 1980, pp. 30-37.
2. Cain, H., and Kraus, B.: Immunofluorescence microscopic demonstration of vimentin filaments in asteroid bodies of sarcoidosis. *Virchows Arch.* 42:213, 1983.
3. Chambers, T. J., Darby, J. A., and Fuller, K.: Mam-  
malian collagenase predisposes bone surfaces to os-  
teoclastic resorption. *Cell Tissue Res.* 241:671,  
1985.
4. Charosky, C. B., Bullough, P. G., and Wilson, P. D.,  
Jr.: Total hip replacement failures: A histological  
evaluation. *J. Bone Joint Surg.* 55A:49, 1973.
5. Chiba, J., Iwaki, Y., Kim, K. J., and Rubash, H. E.:  
The role of cytokines in femoral osteolysis after ce-  
mentless total hip arthroplasty. *Trans. O.R.S.*  
17:350, 1992.
6. Ewald, F. C.: The Knee Society total knee arthro-  
plasty roentgenographic evaluation and scoring sys-  
tem. *Clin. Orthop.* 245:9, 1989.
7. Goldring, S. R., Schiller, A. L., Roelke, M., Rourke,  
C. M., O'Neill, D. A., and Harris, W. H.: The syno-  
vial-like membrane at the bone-cement interface in  
loose total hip replacements and its proposed role in  
bone lysis. *J. Bone Joint Surg.* 65A:575, 1983.
8. Gowens, M., Wood, D. D., Ihrie, E. J., McGuire,  
M. K. B., and Russel, R. G. G.: An interleukin-1-  
like factor stimulates bone resorption in vitro. *Nature*  
306:378, 1983.
9. Heinrich, P. C., Castell, J. V., and Andus, T.: Inter-  
leukin-6 and the acute phase response. *Biochem. J.*  
265:621, 1990.
10. Helle, M., Brakenhoff, J. P. J., and DeGroot, E. R.:  
Interleukin-6 is involved in interleukin-1-induced  
activities. *Eur. J. Immunol.* 18:957, 1988.
11. Howie, D. W., Vernon-Roberts, B., Oakeshott, R.,  
and Manthey, B.: A rat model of resorption of bone  
at the cement-bone interface in the presence of poly-  
ethylene wear particles. *J. Bone Joint Surg.* 70A:257,  
1988.
12. Ishimi, Y., Miyaura, C., and Suda, T.: IL-6 is pro-  
duced by osteoblasts and induces bone resorption. *J.*  
*Immunol.* 145:3297, 1990.
13. Johanson, N. A., Bullough, P. G., Wilson, P. D., Jr.,  
Salvati, E. A., and Ranawat, C. S.: The microscopic  
anatomy of the bone-cement interface in failed total  
hip arthroplasties. *Clin. Orthop.* 218:123, 1987.
14. Jones, S. M., Pinder, I. M., Moran, C. G., and Mal-  
colm, A. J.: Polyethylene wear in uncemented knee  
replacements. *J. Bone Joint Surg.* 74B:18, 1992.
15. Kim, K. J., Rubash, H. E., Wilson, S. C., D'An-  
tonio, J. A., and McClain, E. J.: A histologic and

- biochemical comparison of the interface tissues in cementless and cemented hip prostheses. *Clin. Orthop.* 287:142, 1993.
16. Kurihara, N., Bertolini, D., Suda, T., Akiyama, Y., and Roodman, G. D.: IL-6 stimulates osteoclast-like multinucleated cell formation in long term human marrow cultures by inducing IL-1 release. *J. Immunol.* 144:4226, 1990.
  17. Linder, L., Lindberg, L., and Carlsson, A.: Aseptic loosening of hip prostheses: A histologic and enzyme histochemical study. *Clin. Orthop.* 175:93, 1983.
  18. Lomabardi, A. V., Jr., Mallory, T. H., Vaughn, B. K., and Drouillard, P.: Aseptic loosening in total hip arthroplasty secondary to osteolysis induced by wear debris from titanium-alloy modular femoral heads. *J. Bone Joint Surg.* 71A:1337, 1989.
  19. Maloney, W. J., Jasty, M., Harris, W. H., Galante, J. O., and Callaghan, J. J.: Endosteal erosion in association with stable uncemented femoral components. *J. Bone Joint Surg.* 72A:1025, 1990.
  20. Mirra, J. M., Amstutz, H. C., Matos, M., and Gold, R.: The pathology of the joint tissues in prosthesis failure and its clinical relevance. *Clin. Orthop.* 117:221, 1976.
  21. Nolan, J. F., and Bucknill, T. M.: Aggressive granulomatosis from polyethylene failure in an uncemented knee replacement. *J. Bone Joint Surg.* 74B:23, 1992.
  22. Peters, P., Jr., Engh, G. A., Dwyer, K., and Vinh, T. N.: Osteolysis after total knee arthroplasty without cement. *J. Bone Joint Surg.* 74A:864, 1992.
  23. Schmalzried, T. P., Jasty, M., and Harris, W. H.: Periprosthetic bone loss in total hip arthroplasty. *J. Bone Joint Surg.* 74A:849, 1992.
  24. Sherry, B., and Cerami, A.: Cachectin/tumor necrosis factor exerts endocrine, paracrine, and autocrine control of inflammatory response. *J. Cell. Biol.* 107:1269, 1988.
  25. Simon, P. L., Laydon, J. T., and Lee, J. C.: A modified assay for interleukin-1 (IL-1). *J. Immunol. Methods* 84:85, 1985.
  26. Stern, S. H., and Insall, J. N.: Posterior stabilized prosthesis. *J. Bone Joint Surg.* 74A:980, 1992.
  27. Watanabe, S., Georgescu, H. I., Mendelow, D., and Evans, C. H.: Chondrocyte activation in response to factors produced by a continuous line of lapine synovial fibroblasts. *Exp. Cell Res.* 167:218, 1986.
  28. Willert, H. G., Bertram, H., and Buchhorn, G. H.: Osteolysis in alloarthroplasty of the hip. The role of ultra-high molecular weight polyethylene wear particles. *Clin. Orthop.* 258:95, 1990.

Mi

The  
patie  
care  
of 73  
sive  
perfo  
that  
come  
two  
ing t  
and  
son g  
diti  
secou  
HTO  
ratin  
eithe  
surge  
comp  
nostic  
< 0.0  
tient,  
after  
no pe  
surge  
labor

Pa  
(TKA  
make

From  
Ortho  
School  
timore  
Repr  
Good  
G-1, 50  
Rece

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**